

Aggressive Periodontitis and its Multidisciplinary Focus: Review of the Literature

Periodontitis agresiva y su enfoque multidisciplinario: Revisión de literatura

Maribel Frías-Muñoz DDS¹; Roxana Araujo-Espino DDS, MSc, PhD²;
Víctor Manuel Martínez-Aguilar DDS, MSc³; Teresina Correa Alcalde DDS⁴;
Luis Alejandro Aguilera-Galaviz DDS, MSc, PhD⁴; César Gaitán-Fonseca DDS, MSc, PhD⁴

1. Estudiante, Maestría en Ciencias Biomédicas, Área Ciencias de la Salud, Universidad Autónoma de Zacatecas "Francisco García Salinas" (UAZ), Zacatecas, Zac., Mexico.
2. Docente-investigador, Unidad Académica de Enfermería, Área Ciencias de la Salud, Universidad Autónoma de Zacatecas "Francisco García Salinas" (UAZ), Zacatecas, Zac., Mexico.
3. Docente-investigador, Facultad de Odontología, Universidad Autónoma de Yucatán (UADY), Mérida, Yuc., Mexico.
4. Docente-investigador, Área Ciencias de la Salud, Maestría en Ciencias Biomédicas, Universidad Autónoma de Zacatecas "Francisco García Salinas" (UAZ), Zacatecas, Zac., Mexico.

Correspondence to: Dr. César Gaitán-Fonseca - cgaitan@uaz.edu.mx

Received: 1-V-2017

Accepted: 5-V-2017

Published Online First: 8-V-2017

DOI: <http://dx.doi.org/10.15517/ijds.v0i0.28869>

ABSTRACT

The purpose of this review is to have a current prospect of periodontal diseases and, in particular, aggressive periodontitis. To know its classification and clinical characteristics, such as the extent and age group affected, as well as its distribution in the population, etiology, genetic variations, among other factors that could affect the development of this disease. Also, reference is made to different diagnostic options and, likewise, the current treatment options.

KEYWORDS

Aggressive periodontitis; Periodontal diseases; Diagnosis.

RESUMEN

El propósito de esta revisión es tener un panorama actual de las enfermedades periodontales y, en particular, de la periodontitis agresiva. Conocer su clasificación y características clínicas, como la extensión y grupo etario afectado, así como su distribución en la población, etiología, variaciones genéticas, entre otros factores que pudiesen afectar el desarrollo de dicha enfermedad. Así mismo, se hace referencia a distintas opciones de diagnóstico y, de igual forma, las opciones de tratamiento actuales.

PALABRAS CLAVE

Periodontitis agresiva; Enfermedad periodontal; Diagnóstico.

INTRODUCTION

Periodontal disease is an endogenous microbial disease that damages the dental structure and the periodontium (1). The disease derives from the cellular and humoral response of the host, altering the homeostasis of the periodontal tissues and causing inflammation and destruction by means of bacterial enzymes and virulence factors (2).

Aggressive Periodontitis (AP) is a complex disease that promotes microbial alteration and cellular dysfunction in systemically healthy patients. It begins at any age and prevails in adolescents and young adults. It is characterized by rapid loss of adherence and bone destruction, inconsistent with the amount of microbial deposits present on dental surfaces in local or generalized form (3).

Localized Aggressive Periodontitis (LAP) begins at peri-pubertal age, with interproximal periodontal destruction in primary molar and in no more than two additional affected teeth (4). The presence of dental calcifications on dental surfaces is not frequent; the tissues inflammation and bone-loss patterns are vertical and “U” in form (5). Generalized Aggressive Periodontitis (GAP) affects more than three teeth in addition to the

primary molars and incisors, and it presents loss of interproximal insertion in persons aged <30 years and episodic destruction of alveolar bone (6).

ETIOLOGY

This is a complex disease that possesses four risk factors: subgingival microbiota; individual genetic variations; lifestyle, and systemic factors.

SUBGINGIVAL MICROBIOTA

AP is associated with *Aggregatibacter actinomycetemcomitans*, *Porphyromonas gingivalis*, and abnormal neutrophil function. Both bacteria, due to the action of their virulence factors, act on the host, activating its mechanism of proliferation and destruction, damaging the defense cells and bone resorption, creating periodontal bursae, loss of insertion, mobility, and loss of dental organs (7).

GENETIC VARIABLES

These are associated with biological or endophenotypic intermediaries, which have the potential to modify the host barrier function, inflammatory responses, and microbial colonization patterns. AP comprises a group of distinct conditions, with similar clinical and superimposed presentations

in which each of these is influenced by human genetic variation. A non-protective inflammatory response presents that interacts with the biofilm of the dental surface, generating dysbiotic microbial changes and the establishment of the clinical disease (8).

Hereditary autosomal recessive mechanisms have been related to the appearance and progression of periodontal disease. In AP, there are interindividual differences in the degree of production of inflammatory cytokines, such as Interleukin (IL)-1, Tumor Necrosis Factor alpha (TNF- α), and Prostaglandin E2 (PGE2), following a stimulation due to a leukocytic endotoxin. In addition to the presence of the genetic polymorphism associated with the differences in the interindividual production of IL-1 and TNF- α , there are variants associated with AP, such as the IL-4 gene; (9) this cytokine stimulates the production of B lymphocytes, Immunoglobulin G (IgG), and Immunoglobulin E (IgE) antibodies, and differentiation into T cells inhibits the inflammatory response of the macrophages and the production of IL-1. The polymorphism present in the receptor gene of vitamin D is related with bone density; thus, it is associated with LAP (10).

LIFESTYLE

A person's lifestyle plays a fundamental role in the appearance or inhibition of AP. With respect to the appearance of AP, local and systemic factors are present, such as age, gender, stress, and socioeconomic level. Disease severity increases with age, there is a greater prevalence in women linked with hormonal changes at the pubertal stage, periods of stress diminish the immune response of the organism, and the disease has been associated with low socioeconomic level, which is characterized by deficient habits of hygiene and diet (11).

Stress comprises a risk factor for the development of AP. The effects of the organism's response are anxiety, depression, cognitive alliteration, and an alteration in self-esteem, which give rise to alteration of health behaviors and include negligent oral-hygiene practices and the installation of bruxism. Depression is considered an indicator of stress and is related with the consumption of tobacco, alcohol, and the ingestion of an unhealthy diet, increasing the patient's susceptibility to develop AP. There is individual variability in the immunological response to stress; in some cases, IL-1B increases and regulation of the immune system diminishes. The defense of the immune system against antigens is due to the interaction of the behavior between the innate Central Nervous System (CNS) and the immune system cells. The defense of the immune system against antigens is due to the interaction of the behavior between the CBS and the immune-system cells (12).

Smoking is the most significant risk factor in the prevalence and progression of AP, whose severity depends on the dose consumed (13). The effects of smoking are related with the formation of dentobacterial plaque and the inflammatory response. The physiopathological effects are generated by the nicotine, smoke, and the carbon monoxide resulting from incomplete combustion, which favor a series of molecular events causing the diminution of osteoprotegerin, supporting the interaction of the RANK- RANKL pathway, intervening in the differentiation and activation of osteoclasts, in turn giving rise to resorption (14).

Tobacco smoke increases the proliferation of bacteria in dentobacterial plaque and subgingival microbiota and diminishes the proliferative capacity of the lymphocytes, of Immunoglobulin G2 (IgG2) in the saliva and the synthesis of beta-defensin 2. It generates vasoconstriction and increases platelet

adhesion, augmenting the risk of microvascular occlusion and tissue ischemia reinforced by the CO₂ released, forming carboxyhemoglobin, reducing the contribution of oxygen to the tissue, cellular respiration, and formation of Free Radicals (FR). Nicotine produces an increase in bacterial adhesion to epithelial cells, of the activity of the fibroblastic collagenase enzyme, with a diminution of collagen favoring osteoclastic activity due to the increase of PGE₂. It increases the release of inflammatory cytokines such as IL-6, IL-8, and NTF- α alfa, generating bone loss, destruction of connective tissue, and diminishing repair (15). In addition, it causes peripheral vasoconstriction of the blood vessels and in appearance reduces the clinical signs of gingivitis (16,17).

Ingestion of drugs for the control of systemic diseases such as high blood pressure, diabetes mellitus, and radiotherapy can exert secondary oral effects such as xerostomia, black tongue, recurrent aphthous ulcers, and gingival enlargement, which generate a predisposition for AP. Other drugs, in particular those in liquid or chewable form that contain added sugar, alter the pH and the composition of the plaque. Additionally, drugs such as calcium blockers, Cyclosporine, and anticonvulsants produce gingival hyperplasia (18). These drugs are structurally different, but have in common the action of inhibiting cell calcium uptake, a mechanism implied in the pathogeny of gingival enlargement. It has been demonstrated that young patients exhibit an exaggerated response to drugs, due to the higher level of androgens at the blood level, generating an active metabolite that acts directly on the subpopulations of fibroblasts, this response with greater frequency in the masculine gender (19).

EPIDEMIOLOGY

There are reports in the literature that AP affects 47.2% of the U.S population; the prevalence of AP in adolescents has been

estimated at between 0.1 and 2%. Other studies whose objective was to determine periodontal disease in young population found a prevalence in persons aged between 13 and 20 years of <1%, while in adolescents between the ages of 15 and 17 years, the prevalence was estimated at 0.2% for Caucasians and at 2.6% for Blacks. Similarly, greater prevalence was found in women than in men, and <1% of the population aged under the age of 30 years had AP (20).

The prevalence of LAP in European population varies among adolescents and young adults between 0.1 and 0.2% (21). In industrialized countries, it was found that LAP affects primary dentition in children aged between 5 and 11 years, with a frequency ranging between 0.9 and 4.5% (22,23).

DIAGNOSIS

Periodontal clinical parameters are as follows: Probing Depth (Pd); Level of Clinical Insertion (LCI), and Bleeding on Probing (BOP). Pd is the space that can measure between 1 and 3 mm in the absence of clinical inflammation; a periodontal pocket is defined as the pathological depth of the depth of the periodontal groove, rendered by bone loss and periodontal insertion. For practical clinical effects, a periodontal pocket represents one of the cardinal signs of periodontitis, given that it is produced by the loss of insertion, and can be considered as such from 4mm. The National Informatics Centre (NIC) makes reference to the fibers of the connective gingival tissue that are inserted into the radicular/root cement through the Sharpey fibers. In the clinical ambit, NIC is utilized to refer to the magnitude of support loss, but it depends on the particular radicular length of each tooth.

BOP is the main predictor of periodontal disease and is induced by penetration of the periodontal probe. It should be interpreted in

a global manner, because its presence is not absolutely indicative of disease, while its absence is indeed a reliable indicator of periodontal health.

RADIOGRAPHIC BONE LOSS

Radiographically, periodontal bone pathology presents loss in the continuity of bone and cortical crests, loss of bone height, formation of bone defects, and periodontal ligament enlargement and furcation. Bone loss severity is classified according to the distance from the Cemento-Enamel Junction (CEJ) to the tooth apex: cervical or mild; medium or moderate, or apical or severe.

DIAGNOSTIC ALTERNATIVES

Different complementary diagnostic alternatives to the clinical diagnosis: the use of immunoproteomic approaches implied in the immune response. There is a wide variety of potential proteomic periodontal markers that are included within the immunoproteome: from immunoglobins to bone remodeling proteins. Immunoglobulin M (IgM) is a natural antibody that can bind specific antigens to those to which the host has never been exposed, and it presents traits that allow it to bind to antigens to the degree of invasion, resulting in the activation of the complement as a mechanism of first-line defense, participating in early recognition of bacteria in periodontal disease (24).

C-Reactive Protein (CRP) is a plasma protein that reflects a measurement of the acute-phase response to inflammation, and is one of the markers-of-choice in the follow-up of this response. It is a recognition molecule of patterns that bind specific molecules that are produced during cell death or that are found on the surfaces of diverse bacterial pathogens. The rapid increase of CRP synthesis during the first hours of the progression of an infection suggests its contribution to defense of the host as part of the innate immune response. CRP is produced in response to many types of

distinct lesions of periodontitis, which is found regulated by diverse cytokines. The changes in the cellular and molecular compartments of peripheral blood can be found in patients with periodontitis due to inflammatory changes in the periodontal tissues (25).

TREATMENT

In AP, conventional mechanical therapy and oral hygiene is not sufficient to control the disease. The use of broad-spectrum antibiotics, such as Amoxicillin/clavulanic acid, Metronidazole, Clindamycin, Ciprofloxacin, Tetracycline, and Azithromycin are efficient in pharmacological treatment, in addition to treatments such as surgery, laser therapy, and photodynamic therapy (26).

Photodynamic Therapy (PDT) is a non-invasive tool that functions from the generation of Free Radicals (FR) and oxygen molecules through a photosensitizer placed in an inactive diana tissue, inactivating microorganisms or molecules that react with the light activator. The cytoplasmic membrane of the bacterium is damaged, leading to the inactivation of the membrane transport system, inhibition of the plasma membrane, enzymatic activities, and lipid peroxidation (27), destruction of proteins and ion channels, elimination of critical metabolic enzymes, cell agglutination, and direct inhibition of exogenous virulence factors such as lipopolysaccharides, collagenase, and protease. PDT acts in microorganisms such as fungi, viruses, and protozoans, infections due to simple herpes virus, *P. gingivalis*, *Porphyromonas intermedia*, and *A. actinomycetemcomitans* (28).

REFERENCES

1. Ocampo Yáñez B. R., Marín González M. G. Tratamiento de periodontitis agresiva localizada con plasma rico en plaquetas y aloinjerto óseo. Un caso clínico. *Rev. Odont. Mex.* 2015;19 (2): 106-114.

2. Paknejad M., Khorsand A., Yaghobee S., Motahhari P., Etebarian A., Bayani M., Mehrfard A. Cementogenesis in patients with localized aggressive periodontitis. *J Dent.* 2015;12 (5): 347-351.
3. Aristizábal J. F., Martínez R. Tratamiento ortodóncico y periodontal combinado en pacientes con periodontitis agresiva tratada y controlada. *Rev. Fac. Odontol. Univ. Antioq.* 2014; 26 (1): 180-204.
4. Koss M. A., Castro C. E., Gramajo A. M., López M. E. sIgA, peroxidase and collagenase in saliva of smokers aggressive periodontal patients. *J Oral Biol Craniofac Res.* 2016; 6 (Suppl 1): S24-S28.
5. Merchant S. N., Vovk A., Kalash D., Hovencamp N., Aukhil. Localized aggressive periodontitis treatment response in primary and permanent dentitions. *J. Periodontol.* 2014;85 (12): 1722-1729.
6. Lang N., Bartold P. M., Cullinan M., Jeffcoat M., Mombelli A., Murakami S., Dyke TV. Consensus report: aggressive periodontitis. *Ann. Periodontol.* 1999; 4 (1): 53-53.
7. Ciriaco E. V., Buendía M. D. C. L. Tratamiento multidisciplinario en una paciente con periodontitis agresiva generalizada y diabetes mellitus tipo 1. *Rev. Odont. Mex.* 2014;18 (1): 32-37.
8. Offenbacher S., Divaris K., Barros S. P., Moss K. L., Marchesan J. T., Morelli T., Laudes M. Genome-wide association study of biologically-informed periodontal complex traits offers novel insights into the genetic basis of periodontal disease. *Hum. Mol. Genet.* 2016; 25 (10): 2113-2129.
9. Gutiérrez R. F. Polimorfismo y genética y su relación con la enfermedad periodontal. *Rev. Kiru.* 2015; 5 (2): 127-135.
10. Chatzopoulos G.-S., Doufexi, A.-E., & Kalogirou, F. Association of susceptible genotypes to periodontal disease with the clinical outcome and tooth survival after non-surgical periodontal therapy: A systematic review and meta-analysis. *Med Oral Patol Oral Cir Bucal.* 2016; 21(1): e14–e29.
11. Román R., Zerón A. Factores de riesgo asociados a la enfermedad periodontal. *Rev. Mex. Periodontol.* 2015; 6 (2): 62-66.
12. Ysla-Cheé, R. Estrés y su implicancia en la enfermedad periodontal en escolares. *Rev. Kiru.* 2013;10 (2): 133-139.
13. Bouza-Vera M., Martínez-Abreu J., Carmenate-Rodríguez Y., Betancourt-González M., García-Nicieza M. Pregnancy and oral health. *Rev. Med. Electron.* 2016; 38 (4): 628-634.
14. Elizondo R. T., García MÁQ, Martínez F. L., Palma G. C. El tabaco como factor de riesgo en los tejidos periodontales. *Rev. Mex. Estomatol.* 2015; 2 (3):7-8.
15. Castellanos G. M., Cueto H. M., Boch M., Méndez C. C., Méndez G. L., Castillo F. C. Efectos fisiopatológicos del tabaquismo como factor de riesgo en la enfermedad periodontal. *Rev. Finlay.* 2016; 6 (2): 134-149.
16. Traviesas E. M., Márquez D., Rodríguez R., Rodríguez J., Bordón D. Necesidad del abandono del tabaquismo para la prevención de enfermedad periodontal y otras afecciones. *Rev. Cubana Estomatol.* 2011; 48 (3): 257-267.
17. Martínez-Lima M., Sánchez-Suárez R., Lavandero-Espina A., González del Sol L. Behavior of periodontal emergencies and risk factors in adolescents, Ángel Ameijeiras Polyclinic, Guanabacoa. *Rev. Haban. Cienc. Med.* 2014 ;13 (2):196-206.
18. AlJehani, Y. A. Risk Factors of Periodontal Disease: Review of the Literature. *International J Dent.* 2014, 182513.
19. Campolo G. A., Nuñez C. L., Romero R. P., Rodriguez S. A., Fernandez T. M., Donoso H. F. D. Agrandamiento gingival por ciclosporina: reporte de un caso. *Rev. Clin. Periodoncia Implantol Rehabil. Oral.* 2015; 9 (3): 226-230.
20. Schenkein H. A., Koertge T. E., Sabatini R., Brooks C. N., Gunsolley J. C. Birth weight of infants of mothers with aggressive

- periodontitis. *J. Periodontol.* 2012; 83 (3): 279-286.
21. Renatus A., Trentzsch L., Schönfelder A., Schwarzenberger F., Jentsch H. Evaluation of an Electronic Periodontal Probe Versus a Manual Probe. *Journal of Clinical and Diagnostic Research : JCDR.* 2016; 10 (11): ZH03-ZH07.
 22. Garcia-Madueño N., Gutiérrez-Romero F., Bolaños-Palacios J. Eficacia de la terapia periodontal no quirúrgica en una periodontitis agresiva. *Duazary* 2016; 13 (1): 52-56.
 23. Ardila-Medina C. M., Botero-Zuluaga L., Guzmán-Zuluaga I. C. Comparación de las características sociodemográficas, clínicas y microbiológicas de pacientes con periodontitis agresiva y crónica. *AMC.* 2014; 18 (5): 532-544.
 24. Lin H. S., Talwar H. S., Tarca A. L., Ionan A., Chatterjee M., Ye B., Peshek B. Autoantibody approach for serum-based detection of head and neck cancer. *Cancer Epidemiol.Prev. Biomarkers.* 2007;16 (11): 2396-2405.
 25. Podzimek S., Mysak J., Janatova T., Duskova J. C-Reactive Protein in Peripheral Blood of Patients with Chronic and Aggressive Periodontitis, Gingivitis, and Gingival Recessions. *Mediators of Inflammation.* 2015; 2015: 564858.
 26. Ercan E., Uzun B. C., Ustaoglu G. Effects of Azithromycin versus Metronidazole–Amoxicillin combination as an adjunct to nonsurgical periodontal therapy of generalized aggressive periodontitis. *NJCP.* 2015; 18 (4): 506-510.
 27. Chatzopoulos G.-S., Doufexi A.-E. Photodynamic therapy in the treatment of aggressive periodontitis: A systematic review. *Medicina Oral, Patología Oral y Cirugía Bucal.* 2016; 21 (2): e192-e200.
 28. Cid F., Jara J., Huerta L., Oliva P. Eficacia de terapia fotodinámica como complemento de Terapia convencional periodontal versus terapia convencional en el tratamiento de pacientes adultos con periodontitis crónica: una revisión sistemática con meta análisis. *J. Odontostomatol.* 2016; 10 (2): 315-323.



Attribution (BY-NC) - (BY) You must give appropriate credit, provide a link to the license, and indicate if changes were made. You may do so in any reasonable manner, but not in any way that suggest the licensor endorses you or your use. (NC) You may not use the material for commercial purposes.